

CAS 1163-19-5

Substance name 2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (also known as BDE-209)

Toxicity

BDE -209 is the primary congener found in Deca-BDE. Thyroid and liver appear to be the most sensitive tissues to toxicity of Deca-BDE in animal studies. In 2008, EPA determined that Deca-BDE had “suggestive evidence of carcinogenic potential” in humans.¹ This is based on liver tumors in rats and male mice, and thyroid gland follicular cell hyperplasia and thyroid tumors in male mice in oral studies with Deca-BDE.^{1,2} Rats and mice exposed to Deca-BDE in their early postnatal period, were observed to have neurodevelopmental effects as they matured.^{3,4,5,6} In the environment, BDE-209 is likely to degrade into less-brominated, more toxic BDEs.⁷

Exposure

Deca BDE is listed as a Persistent, Bioaccumulative and Toxic (PBT) chemical under Washington State's PBT rule (WAC 173-333-320).⁷ This chemical is widely used as a flame retardant in high impact polystyrene and other polymers, in coatings and adhesive systems such as the back coatings for carpets, and in non-clothing textiles.^{1,7} It is a high production volume chemical that has not been reported directly in children's products but has been found in indoor air and dust and in biomonitoring studies.^{1,7,8,9}

References

1. U.S. Environmental Protection Agency. Toxicological review of Decabromodiphenyl Ether (BDE-209), June 2008. <http://www.epa.gov/iris/toxreviews/0035tr.pdf>.
2. National Toxicology Program (1986) Toxicology and carcinogenesis studies of decabromodiphenyl oxide (CAS No. 1163-19-5) in F344/N rats and B6C3F1 mice (feed studies). Public Health Service, U.S. DHHS; NTP TR 309. http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr309.pdf.
3. Viberg, H; Fredriksson, A; Eriksson, P. (2007) Changes in spontaneous behavior and altered response to nicotine in the adult rat, after neonatal exposure to the brominated flame retardant, decabrominated diphenyl ether (PBDE 209). *Neurotoxicol* 28:136–142.
4. Rice, DC; Reeve, EA; Herlihy, A; et al. (2007) Developmental delays and locomotor activity in the C57BL6/J mouse following neonatal exposure to the fully-brominated PBDE, decabromodiphenyl ether. *Neurotoxicol Teratol* 29:511–520.
5. Viberg, H; Fredriksson, A; Jakobsson, E; et al. (2003) Neurobehavioral derangements in adult mice receiving decabromodiphenyl ether (PBDE 209) during a defined period of neonatal brain development. *Toxicol Sci* 76:112–120.
6. Darnerud et al. (2001) Polybrominated diphenyl ethers: occurrence, dietary exposure and toxicology. *Environmental Health Perspectives* 109(Supplement 1): 49-68.
7. WA Departments of Ecology and Health. Washington State Polybrominated Diphenyl Ether (PBDE) Chemical Action Plan. January 2006. <http://www.ecy.wa.gov/biblio/0507048.html>.
8. Schecter et al., 2003. Polybrominated diphenyl ethers (PBDEs) in U.S. mother's milk. *Environmental Health Perspectives* 111(14): 1723-1729.
9. Butt et al., 2004. Spatial distribution of polybrominated diphenyl ethers in southern Ontario as measured in indoor and outdoor window organic films. *Environmental Science & Technology* 38(3):724-731.